

Addition and multiplication of beta-expansions in generalized Tribonacci base

P. AMBROŽ

*Department of Mathematics, Faculty of Nuclear Science and Physical Engineering,
Czech Technical University Trojanova 13, 120 00 Praha 2, Czech Republic*

and

*LIAFA, CNRS UMR 7089 2 place Jussieu,
752 51 Paris Cedex 05, France
e-mail: ampy@linux.fjfi.cvut.cz*

Z. MASÁKOVÁ

*Department of Mathematics, Faculty of Nuclear Science and Physical Engineering,
Czech Technical University Trojanova 13, 120 00 Praha 2, Czech Republic*

e-mail: masakova@km1.fjfi.cvut.cz

and

E. PELANTOVÁ

*Department of Mathematics, Faculty of Nuclear Science and Physical Engineering,
Czech Technical University Trojanova 13, 120 00 Praha 2, Czech Republic*

e-mail: pelantova@km1.fjfi.cvut.cz

ABSTRACT

We study properties of β -numeration systems, where $\beta > 1$ is the real root of the polynomial $x^3 - mx^2 - x - 1$, $m \in \mathbb{N}$, $m \geq 1$. We consider arithmetic operations on the set of β -integers, i.e. on the set of numbers whose greedy expansion in base β has no fractional part. We show that the number of fractional digits arising under addition of β -integers is at most 5 for $m \geq 3$ and 6 for $m = 2$, whereas under multiplication it is at most 6 for all $m \geq 2$. We thus generalize the results known for Tribonacci numeration system, i.e. for $m = 1$. We summarize the combinatorial properties of infinite words naturally defined by β -integers. We point out the differences between the structure of β -integers in cases $m = 1$ and $m \geq 2$.

Keywords: beta-expansion, numeration system, Pisot number.

1. Introduction

In this paper we study the properties of a non-standard numeration system related to an irrational base β for a class of cubic numbers β .

The definition of a numeration system based on a real number $\beta > 1$ is given by the fact that every non-negative x has a unique expression in the form of its β -expansion,

$$x = \sum_{i=-\infty}^k a_i \beta^i,$$

where the coefficients a_i are non-negative integers and

$$0 \leq x - \sum_{i=n}^k a_i \beta^i < \beta^n \quad \text{for all } n \in \mathbb{Z}, n \leq k. \quad (1)$$

A real number x for which the β -expansion of $|x|$ is of the form $|x| = \sum_{i=0}^k a_i \beta^i$ is called a β -integer. The set of β -integers is denoted by \mathbb{Z}_β . We say that a real x has a finite β -expansion, if x is an element of the set

$$\text{Fin}(\beta) := \bigcup_{n=0}^{\infty} \frac{1}{\beta^n} \mathbb{Z}_\beta.$$

If $\beta = 10$, then the β -expansion of x is the ordinary decimal expansion of x and $\mathbb{Z}_\beta = \mathbb{Z}$. An equality $\mathbb{Z}_\beta = \mathbb{Z}$ holds for every rational integer $\beta \in \mathbb{Z}$, $\beta > 1$.

If $\beta \notin \mathbb{Z}$, the condition (1) implies that the coefficients a_i of the β -expansion satisfy $a_i \in \{0, 1, \dots, \lfloor \beta \rfloor\}$ for all $i \leq k$. However, not all power series with such coefficients corresponds to a β -expansion of some real x . For the characterization of admissible sequences of coefficients one needs to introduce the so-called Rényi expansion of 1,

$$d_\beta(1) = t_1 t_2 t_3 \cdots, \quad \text{where } t_1 = \lfloor \beta \rfloor \text{ and}$$

$$\sum_{n=2}^{\infty} \frac{t_n}{\beta^n} \text{ is the } \beta\text{-expansion of } 1 - \frac{t_1}{\beta}.$$

A number β , for which $d_\beta(1)$ is eventually periodic is called a Parry number. If moreover $d_\beta(1) = t_1 t_2 t_3 \cdots$ has only a finite number of non-zero entries t_i , then β is called a simple Parry number and the vanishing coefficients at the end of $d_\beta(1)$ are omitted. Expansions of β -integers are characterized by the Parry condition [18]:

$\sum_{i=0}^k a_i \beta^i$ with $a_i \in \{0, 1, 2, \dots\}$ is a β -expansion of some $x > 0$ if and only if $a_i a_{i-1} \cdots a_0 \prec t_1 t_2 t_3 \cdots$, i.e. $a_i a_{i-1} \cdots a_0$ is lexicographically strictly smaller than $t_1 t_2 t_3 \cdots$ for any $i = k, k-1, \dots, 0$.

Unlike numeration systems with an integer base, the set \mathbb{Z}_β of β -integers is not closed under addition and multiplication if $\beta \notin \mathbb{Z}$. However, this does not exclude that $\text{Fin}(\beta)$ is a ring. A necessary condition so that $\text{Fin}(\beta)$ is closed under addition and multiplication is that β is a Pisot-Vijayaraghavan (PV) number and a simple Parry number [1, 12]. Recall that PV numbers are defined as real algebraic integers greater than 1 such that their conjugates are inside the unit complex circle. It is known that every PV number is a Parry number.

Some sufficient conditions on a simple Parry number β so that $\text{Fin}(\beta)$ is a ring are also known¹. In [12] it is shown that if $d_\beta(1) = t_1 t_2 \cdots t_m$ satisfies $t_1 \geq t_2 \geq \cdots \geq$

¹We say that the number β has Property (F) if the set $\text{Fin}(\beta)$ is a ring

$t_m > 0$ then $\text{Fin}(\beta)$ is a ring. Hollander [15] has shown that $t_1 > t_2 + t_3 + \dots + t_m$ is a sufficient condition so that $\text{Fin}(\beta)$ is a ring. For cubic PV units β Akiyama [1] has shown that $\text{Fin}(\beta)$ is a ring if and only if β is a simple Parry number, i.e. $d_\beta(1)$ is finite. A weaker form of the condition (F) is the so-called weak finiteness property, studied in [4].

For performing the arithmetic operations in the numeration system based on β it is important not only to know that the result has a finite fractional part, but to know the length of the fractional part. We therefore study the quantities

$$L_\oplus = L_\oplus(\beta) := \min\{n \in \mathbb{N}_0 \mid \forall x, y \in \mathbb{Z}_\beta, x + y \in \text{Fin}(\beta) \Rightarrow x + y \in \beta^{-n}\mathbb{Z}_\beta\},$$

$$L_\otimes = L_\otimes(\beta) := \min\{n \in \mathbb{N}_0 \mid \forall x, y \in \mathbb{Z}_\beta, xy \in \text{Fin}(\beta) \Rightarrow xy \in \beta^{-n}\mathbb{Z}_\beta\}.$$

In [2] it is shown that for β a PV number, one has $L_\oplus < +\infty$. In [14] the same result is given for L_\otimes . First known values for L_\oplus, L_\otimes have been provided in [9] for quadratic PV units. Article [14] states exact values and estimates on L_\oplus, L_\otimes for arbitrary quadratic PV numbers. The first cubic case was studied by Messaoudi. In [16] he shows for the so-called Tribonacci numeration system, i.e. for the base β with minimal polynomial $x^3 - x^2 - x - 1$, that $L_\otimes \leq 9$, and states the conjecture of Arnoux that $L_\otimes = 3$. In [17] he improves the upper bound to $L_\otimes \leq 6$. In [6] it is shown that $4 \leq L_\otimes \leq 5$ and $5 \leq L_\oplus \leq 6$. Final exact value $L_\oplus = 5$ has been given with an elegant proof by Bernat [8].

In this paper we study a class of cubic units $\beta > 1$, roots of

$$x^3 = mx^2 + x + 1, \quad m \in \mathbb{N}.$$

We call such base β the generalized Tribonacci number, since the Tribonacci case is obtained for $m = 1$. The equation $x^3 = mx^2 + x + 1$ has a unique real solution β . It satisfies $m < \beta < m + 1$. The other two roots β' and β'' are mutually complex conjugates $|\beta'| = |\beta''| < 1$. Obviously, we have

$$\beta\beta'\beta'' = 1, \quad \beta + \beta' + \beta'' = m.$$

The Rényi expansion of 1 is $d_\beta(1) = m11$. Therefore the digits in a β -expansion take values in $\{0, 1, \dots, m\}$ and the Parry condition implies for $a_0, a_1, \dots, a_k \in \mathbb{N}_0$ that

$$\sum_{i=0}^k a_i \beta^i \text{ is a } \beta\text{-expansion} \iff a_i a_{i-1} a_{i-2} < m11 \quad \text{for all } i = 2, 3, \dots, k.$$

Due to the results of Frougny and Solomyak [12] or Akiyama [2] the set $\text{Fin}(\beta)$ is a ring for all m .

Our aim is to provide estimates on the quantities $L_\oplus(\beta), L_\otimes(\beta)$ for these generalized Tribonacci numbers β . Since the Tribonacci case $m = 1$ has been already solved, we consider $m \geq 2$.

Lower bounds on $L_\oplus(\beta), L_\otimes(\beta)$ can be clearly obtained as the length of the fractional part of the sum, respectively product of some elements $x, y \in \mathbb{Z}_\beta$.

Proposition 1.1 *Let β be the real root of $x^3 = mx^2 + x + 1$. Then*

$$L_{\oplus}(\beta) \geq \begin{cases} 5 & \text{for } m = 2, \\ 4 & \text{for } m \geq 3, \end{cases} \quad L_{\otimes}(\beta) \geq 4, \quad \text{for } m \geq 2.$$

Proof. We have for $m = 2$,

$$(\beta^4 + \beta^2 + 2\beta) - (\beta^3 + 2) = 2\beta^3 + 2 + \beta^{-3} + \beta^{-4} + \beta^{-5},$$

$$2 \times (\beta^6 + 2\beta^5 + 2\beta^3 + \beta^2 + 2) = \beta^7 + \beta^6 + 2\beta^4 + 2\beta + \beta^{-2} + 2\beta^{-3} + \beta^{-4},$$

and for $m \geq 3$,

$$\begin{aligned} (m\beta^3 + m) + (m\beta^3 + m) &= 2 \times (m\beta^3 + m) = \\ &= \beta^4 + (m-1)\beta^3 + (m-1)\beta^2 + 2\beta + \\ &\quad + (m-3)\beta^{-1} + (m-1)\beta^{-2} + 2\beta^{-3} + \beta^{-4}. \end{aligned}$$

□

The main result of this paper is providing the upper bounds on $L_{\oplus}(\beta)$ and $L_{\otimes}(\beta)$. Using a specific method of estimation, which will be explained in the next section, we obtain $5 \leq L_{\oplus} \leq 6$, for $m = 2$, and $4 \leq L_{\oplus} \leq 5$, for $m \geq 3$, (Theorem 3.1). Let us mention that for two special m the exact value of $L_{\oplus}(\beta)$ is known [8], namely for $m = 2$ is $L_{\oplus}(\beta) = 5$ and for $m = 3$ is $L_{\oplus}(\beta) = 4$. Further, we study $L_{\otimes}(\beta)$. We establish $4 \leq L_{\otimes} \leq 6$ for $m \geq 2$, (Theorem 3.2). Note that the exact value of $L_{\otimes}(\beta)$ is not known even in Tribonacci case $m = 1$.

The proofs of Theorems 3.1 and 3.2 use real quadratic forms and their differences. It is noteworthy that the similar computation was the bottleneck of [5] and [13] in studying topology of Thurston-Rauzy fractals associated with β -expansions.

Although we study the numeration systems mainly from the arithmetical point of view, its combinatorial properties are also very interesting. At the end of the paper we present some combinatorial properties of the set of β -integers which can be derived from general results in the literature for the class of generalized Tribonacci numbers.

2. Methods of upper estimation of $L_{\oplus}(\beta)$, $L_{\otimes}(\beta)$

Let us present the methods for finding upper estimates of L_{\oplus} , L_{\otimes} and the one chosen for the class of generalized Tribonacci numbers explain in detail. Two methods for estimation of L_{\oplus} , L_{\otimes} are known. First of them uses the so-called Meyer property of the set of β -integers, namely that

$$\mathbb{Z}_{\beta} + \mathbb{Z}_{\beta} \subset \mathbb{Z}_{\beta} + F, \quad \text{for a finite set } F.$$

In [3] one studies the question of minimizing the finite set F . For multiplication, one has a similar property [14]: If β is a PV number, then

$$\mathbb{Z}_{\beta} \cdot \mathbb{Z}_{\beta} \subset \mathbb{Z}_{\beta} + G, \quad \text{for a finite set } G.$$

This method is used in [6] to find values of $L_{\oplus}(\beta)$, $L_{\otimes}(\beta)$ where β is a PV number, solution of the equation $x^3 = 25x^2 + 15x + 2$.

The second much more used method for estimation of L_{\oplus} , L_{\otimes} is based on the following theorem. Several versions of this method are employed in [6, 8, 14, 16, 17].

Theorem 2.1 ([14]) *Let $\beta > 1$ be an algebraic number, and let β' be its conjugate. For $z \in \mathbb{Q}(\beta)$ we denote by z' the image of z under the field isomorphism $' : \mathbb{Q}(\beta) \rightarrow \mathbb{Q}(\beta')$. If*

$$H := \sup\{|z'| \mid z \in \mathbb{Z}_{\beta}\} < +\infty,$$

$$K := \inf\{|z'| \mid z \in \mathbb{Z}_{\beta} \setminus \beta\mathbb{Z}_{\beta}\} > 0,$$

then

$$\left(\frac{1}{|\beta'}\right)^{L_{\oplus}} < \frac{2H}{K} \quad \text{and} \quad \left(\frac{1}{|\beta'}\right)^{L_{\otimes}} < \frac{H^2}{K}.$$

In the above theorem we require existence of at least one conjugate of β such that the constant H is finite and K is positive. The former is ensured if $|\beta'| < 1$. To decide whether $K > 0$ or $K = 0$ is much more complicated. In the paper [6] there is given a sufficient condition for a number β so that $K = 0$. An example for such number β is the already mentioned solution of the equation $x^3 = 25x^2 + 15x + 2$. Indeed, for such β one cannot employ Theorem 2.1. A sufficient condition for β so that $K > 0$ was given by Akiyama [1]. Akiyama proved that for β PV unit such that $\text{Fin}(\beta)$ is a ring, the origin is an inner point of the central tile in the conjugated plane, i.e. of the closure of the set $\{z' \mid z \in \mathbb{Z}_{\beta}\}$. This implies K is positive for all conjugates of β . In such a case one chooses the conjugate which provides the best estimates on L_{\oplus} , L_{\otimes} .

Our aim is to provide estimates on the quantities $L_{\oplus}(\beta)$, $L_{\otimes}(\beta)$ for the generalized Tribonacci numbers β . We use the method based on Theorem 2.1. Recall that one has to choose a suitable conjugate of β . However, in the case of generalized Tribonacci numbers, both conjugates of β have the same modulus < 1 , and thus both of them will provide the same values of constants H, K . Moreover, since β is a cubic PV unit with finite Rényi expansion of 1, set $\text{Fin}(\beta)$ is a ring and therefore constant K in Theorem 2.1 is positive.

It is a difficult problem to determine the exact values of H and K . However, in order to obtain bounds on L_{\oplus} and L_{\otimes} it suffices to find some “reasonable” upper estimate on H and lower estimate on K .

Upper estimate on H . For a given $p \in \mathbb{N}$, every $z \in \mathbb{Z}_{\beta}$ can be written as

$$z = (a_0 + a_1\beta + \dots + a_{p-1}\beta^{p-1}) + \beta^p(a_p + \dots + a_{2p-1}\beta^{p-1}) + \dots,$$

with the integer coefficients satisfying the admissibility condition $a_i a_{i-1} a_{i-2} \prec m11$. Using triangular inequality, the modulus of the conjugate z' can be estimated by

$$|z'| \leq (1 + |\beta'|^p + |\beta'|^{2p} + \dots) \text{Max}_p = \frac{\text{Max}_p}{1 - |\beta'|^p} =: H_p, \quad (2)$$

where

$$\text{Max}_p := \max\{|z'| \mid z \in \mathbb{Z}_\beta, |z| < \beta^p\}. \quad (3)$$

We therefore have

$$H \leq H_p, \quad \text{for all } p \in \mathbb{N}.$$

Lower estimate on K . For a given $p \in \mathbb{N}$, and for all $z \in \mathbb{Z}_\beta$ we have

$$|z'| \geq |a_0 + a_1\beta + \cdots + a_{p-1}\beta^{p-1}| - |\beta'|^p |a_p + a_{p+1}\beta' + \cdots|.$$

Note that the last term on the right hand side is a modulus of a β -expansion of a conjugate of some $x \in \mathbb{Z}_\beta$ and therefore it is bounded by H (or H_q for some $q \in \mathbb{N}$).

With the notation

$$\text{Min}_p := \min\{|z'| \mid z \in \mathbb{Z}_\beta \setminus \beta\mathbb{Z}_\beta, |z| < \beta^p\}, \quad (4)$$

we obtain an estimate

$$K \geq K_{p,q} := \text{Min}_p - |\beta'|^p H_q, \quad \text{for all } p, q \in \mathbb{N}. \quad (5)$$

We see that the problem of the upper bounds on L_\oplus and L_\otimes has been reduced to the problem of finding Max_p and Min_p for a suitable value of p . Recall from the definition of Max_p and Min_p (equations (3) and (4)) that both of them are calculated using the modulus $|z'|$ of the image of β -integers $|z| < \beta^p$ under the field morphism. It turns more convenient to study the square of this quantity.

Let us consider a β -integer z , such that $|z| < \beta^p$, i.e. z is of the form $z = a_0 + a_1\beta + a_2\beta^2 + \cdots + a_{p-1}\beta^{p-1}$, where the integer coefficients a_i satisfy the admissibility condition $a_i a_{i-1} a_{i-2} \prec m11$. Since the conjugates of β are mutually complex conjugated we have

$$|z'|^2 = z'z'' = \sum_{i=0}^{p-1} a_i \beta'^i \sum_{j=0}^{p-1} a_j \beta''^j = \sum_{i=0}^{p-1} \sum_{j=0}^{p-1} a_i a_j \beta'^i \beta''^j.$$

We use the relation $\beta\beta'\beta'' = 1$ to derive for $j > i$,

$$a_i a_j \beta'^i \beta''^j + a_j a_i \beta'^j \beta''^i = a_i a_j \beta'^{j-i} \frac{1}{\beta^i} + a_i a_j \beta''^{j-i} \frac{1}{\beta^i} = \frac{a_i a_j}{\beta^i} 2\Re(\beta'^{j-i}).$$

Substituting this to the expression for $|z'|^2$, we obtain

$$|z'|^2 = \sum_{k=0}^{p-1} \frac{a_k^2}{\beta^k} + \sum_{0 \leq i < j \leq p-1} \frac{a_i a_j}{\beta^i} 2\Re(\beta'^{j-i}). \quad (6)$$

Therefore, $|z'|^2$ is a real quadratic form of integer variables a_0, a_1, \dots, a_{p-1} . In order to express it in a simpler form, we need to determine the coefficients $2\Re(\beta'^{j-i})$. Let us denote $c_k := 2\Re(\beta'^k)$. For calculation of c_k we find a recurrent formula,

$$\begin{aligned} c_k c_1 &= (\beta'^k + \beta''^k)(\beta' + \beta'') = \beta'^{k+1} + \beta'^k \beta'' + \beta''^k \beta' + \beta''^{k+1} \\ &= c_{k+1} + \beta' \beta'' (\beta'^{k-1} + \beta''^{k-1}) \\ &= c_{k+1} + \frac{1}{\beta} c_{k-1}. \end{aligned}$$

Let us enumerate several initial coefficients,

$$c_0 = 2, \quad (7a)$$

$$c_1 = -\left(\frac{1}{\beta} + \frac{1}{\beta^2}\right), \quad (7b)$$

$$c_2 = -\left(\frac{2}{\beta} - \frac{1}{\beta^2} - \frac{2}{\beta^3} - \frac{1}{\beta^4}\right), \quad (7c)$$

$$c_3 = \frac{3}{\beta^2} + \frac{2}{\beta^3} - \frac{3}{\beta^4} - \frac{3}{\beta^5} - \frac{1}{\beta^6}, \quad (7d)$$

$$c_4 = \frac{2}{\beta^2} - \frac{4}{\beta^3} - \frac{7}{\beta^4} + \frac{6}{\beta^6} + \frac{4}{\beta^7} + \frac{1}{\beta^8}. \quad (7e)$$

Note that for $m \geq 3$ the coefficients satisfy $c_1 < 0$, $c_2 < 0$, $c_3 > 0$ and $c_4 > 0$.

Our aim is to find Max_p and Min_q for suitable p, q so that the bounds on L_{\oplus} , L_{\otimes} obtained by Theorem 2.1 are best possible. Computer experiments show that optimal constants are H_3 , $K_{5,3}$, that is to say the values of H_n and $K_{m,n}$ change too little with increasing n, m and do not provide better estimates on the values of L_{\oplus} and L_{\otimes} . Thus we need to calculate Max_3 and Min_5 .

Since we are interested in Max_3 , the real quadratic form $|z'|^2$ to be examined simplifies to

$$Q_1(a_2, a_1, a_0) := |z'|^2 = a_0^2 + \frac{a_1^2}{\beta} + \frac{a_2^2}{\beta^2} + a_0 a_1 c_1 + a_0 a_2 c_2 + \frac{a_1 a_2}{\beta} c_1. \quad (8)$$

In order to calculate the value of Max_3 we have to find the maximum of the quadratic form $Q_1(a_2, a_1, a_0)$ on the set

$$\mathcal{S} := \{(a_2, a_1, a_0) \in \mathbb{Z}^3 \mid 0 \leq a_i \leq m, a_2 a_1 a_0 < m11\}. \quad (9)$$

Similarly, for calculation of Min_5 , we use the quadratic form

$$\begin{aligned} Q_2(a_4, a_3, a_2, a_1, a_0) := |z'|^2 &= a_0^2 + \frac{a_1^2}{\beta} + \frac{a_2^2}{\beta^2} + \frac{a_3^2}{\beta^3} + \frac{a_4^2}{\beta^4} + a_0 a_1 c_1 + a_0 a_2 c_2 + \\ &+ a_0 a_3 c_3 + a_0 a_4 c_4 + \frac{a_1 a_2}{\beta} c_1 + \frac{a_1 a_3}{\beta} c_2 + \frac{a_1 a_4}{\beta} c_3 + \frac{a_2 a_3}{\beta^2} c_1 + \frac{a_2 a_4}{\beta^2} c_2 + \frac{a_3 a_4}{\beta^3} c_1. \end{aligned} \quad (10)$$

We have to find a minimum of this quadratic form on the set

$$\mathcal{S} := \{(a_4, a_3, a_2, a_1, a_0) \in \mathbb{Z}^5 \mid m \geq a_i \geq 0, a_0 > 0 \text{ and } a_{k+2} a_{k+1} a_k < m11\}. \quad (11)$$

The condition $a_0 \neq 0$ in the definition of set \mathcal{S} corresponds to the condition $z \in \mathbb{Z}_{\beta} \setminus \beta \mathbb{Z}_{\beta}$ in the definition of Min_p .

For finding the extremal values of the quadratic forms, we inspect the first differences of the quadratic form for each variable a_i , each time fixing all yet appointed values of the variables. The difference in a variable a_i will be denoted

$$\Delta_{a_i} := Q(\dots, a_i + 1, \dots) - Q(\dots, a_i, \dots),$$

where $a_i \leq m - 1$ and the variables a_j , $j \neq i$ take values at most m . Using this method we further prove the following result.

Proposition 2.2 *Let $\beta > 1$ be the real root of the equation $x^3 = mx^2 + x + 1$ for $m \in \mathbb{N}$, $m \geq 2$. Then*

$$\begin{aligned} \text{Max}_3 &= \max\{|z'| \mid z \in \mathbb{Z}_\beta, |z| < \beta^3\} = m, \\ \text{Min}_5 &= \min\{|z'| \mid z \in \mathbb{Z}_\beta \setminus \beta\mathbb{Z}_\beta, |z| < \beta^5\} = \begin{cases} \beta^{-2}(\beta + 1) & \text{for } m \geq 3 \\ |1 + 2\beta'^2 + 2\beta'^4| & \text{for } m = 2. \end{cases} \end{aligned}$$

Since the proof of this proposition mainly consists of a rather technical analysis of the behavior of the quadratic forms Q_1, Q_2 , we postpone it to a separate Section 4.

3. The main result

In the preceding sections we have found the values of Max_3 and Min_5 . According to the definitions (2) and (5) we have for $m \geq 3$,

$$H_3 = \frac{\text{Max}_3}{1 - |\beta'|^3} = \frac{m\beta^{3/2}}{\beta^{3/2} - 1}$$

and

$$K_{5,3} = \text{Min}_5 - |\beta'|^5 H_3 = \frac{\beta + 1}{\beta^2} - \frac{m}{\beta(\beta^{3/2} - 1)}.$$

Now, we will use these values to obtain upper estimates on the value of L_\oplus .

Theorem 3.1 *Let $\beta > 1$ be the real root of the equation $x^3 = mx^2 + x + 1$ for $m \in \mathbb{N}$. Then*

$$\begin{aligned} 4 &\leq L_\oplus(\beta) \leq 5 && \text{for } m \geq 3, \\ 5 &\leq L_\oplus(\beta) \leq 6 && \text{for } m = 2. \end{aligned}$$

Proof. The upper bound $L_\oplus \leq 6$ for $m = 2$ can be easily checked numerically. Since

$$\frac{2H}{K} < \frac{2H_3}{K_{5,3}} \simeq 18.4596 < \left(\frac{1}{|\beta'|}\right)^7 \simeq 26.3628,$$

according to the Theorem 2.1 we have $L_\oplus \leq 6$.

Let us assume $m \geq 3$. Again according to the Theorem 2.1 we have

$$\left(\frac{1}{|\beta'|}\right)^{L_\oplus} < \frac{2H}{K} \leq \frac{2H_3}{K_{5,3}}.$$

We will prove the inequality

$$\frac{2H_3}{K_{5,3}} = \frac{2m\beta^{3/2}}{\beta^{3/2} - 1} \frac{1}{\frac{\beta+1}{\beta^2} - \frac{m}{\beta(\beta^{3/2}-1)}} < \beta^3 = \frac{1}{|\beta'|^6}, \quad (12)$$

which implies $L_\oplus \leq 5$.

After a few simple operations the inequality (12) is transformed into

$$2m\beta^{1/2} + m\beta < (\beta + 1)(\beta^{3/2} - 1). \quad (13)$$

For $m \geq 4$ the proof of the inequality (13) will be based on the fact that $\beta > \beta^{1/2} + 2$, which holds since

$$\beta > \beta^{1/2} + 2 \Leftrightarrow (\beta - 2)^2 > \beta \Leftrightarrow m\beta + 1 + \frac{1}{\beta} - 4\beta + 4 > \beta.$$

Now we estimate the left hand side of (13)

$$\begin{aligned} m\beta^{1/2}(2 + \beta^{1/2}) &< \beta^{1/2}m\beta < \beta^{1/2} \left(m\beta + \frac{1}{\beta} \right) = \beta^{1/2}(\beta^2 - 1) = \\ &= \beta^{1/2}(\beta - 1)(\beta + 1) = (\beta + 1)(\beta^{3/2} - \beta^{1/2}) < (\beta + 1)(\beta^{3/2} - 1), \end{aligned}$$

which proves (13).

In the omitted case $m = 3$, the inequality (13) can be proved directly. \square

The following theorem yields the main result for multiplication – the estimates of the value of the coefficient L_{\otimes} .

Theorem 3.2 *Let $\beta > 1$ be the real root of the equation $x^3 = mx^2 + x + 1$ for $m \in \mathbb{N}$, $m \geq 2$. Then*

$$4 \leq L_{\otimes}(\beta) \leq 6.$$

Proof. Let us assume $m \geq 3$. According to the Theorem 2.1 it suffices to show

$$\frac{H^2}{K} \leq \frac{H_3^2}{K_{5,3}} = \frac{m2}{(1 - \beta^{-3/2})^2} \frac{1}{\frac{\beta+1}{\beta^2} - \beta^{-5/2} \frac{m}{1 - \beta^{-3/2}}} < \frac{1}{|\beta'|^7},$$

which can be transformed into

$$(1 + m)\beta^3 + \beta^2 + (m + 2)\beta + 1 > ((m + 2)\beta + m^2 + 2) \beta^{3/2}. \quad (14)$$

By $m \leq \beta < m + 1$ we can estimate the left hand side of (14) as

$$\begin{aligned} (1 + m)\beta^3 + \beta^2 + (m + 2)\beta + 1 &\geq \\ &(1 + m)m^3 + m^2 + (m + 2)m + 1 \geq m^3(1 + m) + m^2, \end{aligned}$$

whereas the right hand side as

$$\begin{aligned} ((m + 2)\beta + m^2 + 2) \beta^{3/2} &< \\ &((m + 2)(m + 1) + m^2 + 2)(m + 1)^{3/2} \leq 3m^2(m + 1)^{3/2}, \end{aligned}$$

where the last inequality holds for $m \geq 4$. The omitted cases will be treated separately at the end of the proof.

Using last two estimates we have

$$3m^2(m + 1)^{3/2} \leq m^2 + m^3(1 + m).$$

It is easy to check that this inequality holds for $m \geq 10$.

Since our estimates were too rough to check the validity of (14) for $m \leq 9$, we verified the proposition numerically for these cases as well as for the case $m = 2$ which was omitted at the beginning of the proof. \square

Let us mention that computer experiments support the hypothesis that $L_{\otimes}(\beta) = 4$ for $m \geq 2$.

4. Proof of Proposition 2.2

In this section we provide the proof of Proposition 2.2. For that we will use the notions of quadratic forms introduced in Section 2. We start with determining the value of Min_5 , since the results will be used in the other part of the proof for Max_3 .

Determining Min_5 . In the case $m = 2$ the set \mathcal{S}_2 defined in (11) has only 79 elements. It is easy to enumerate them one by one and to find the value of Min_5 . Therefore, from now we will consider $m \geq 3$.

1) Difference Δ_{a_0} . The inspected first difference of the quadratic form Q_2 is

$$\Delta_{a_0} = 2a_0 + 1 + c_1a_1 + c_2a_2 + c_3a_3 + c_4a_4.$$

Since $a_0 \geq 1$, $c_1 < 0$, $c_2 < 0$, $c_3 > 0$, $c_4 > 0$ and $a_1, \dots, a_4 \in \{0, \dots, m\}$ we have

$$\begin{aligned} \Delta_{a_0} &\geq 3 + mc_1 + mc_2 = 3 - m \left(\frac{1}{\beta} + \frac{1}{\beta^2} \right) - m \left(\frac{2}{\beta} - \frac{1}{\beta^2} - \frac{2}{\beta^3} - \frac{1}{\beta^4} \right) \\ &\geq 3 - \frac{3m}{\beta} > 0, \end{aligned}$$

where the last inequality follows from the fact that $m = \lfloor \beta \rfloor$.

Therefore, the form is increasing in the variable a_0 and so the minimum is reached at the smallest possible value of a_0 , i.e. at $a_0 = 1$. From now on we will consider $a_0 = 1$.

2) Difference Δ_{a_1} . The inspected difference is

$$\Delta_{a_1} = \frac{2a_1 + 1}{\beta} + c_1 + c_1 \frac{a_2}{\beta} + c_2 \frac{a_3}{\beta} + c_3 \frac{a_4}{\beta}.$$

Since $c_1 < 0$, $c_2 < 0$, $c_3 > 0$ we have

$$\Delta_{a_1} \geq \frac{2a_1}{\beta} - \frac{1}{\beta^2} - \frac{m}{\beta} \left(\frac{1}{\beta} + \frac{1}{\beta^2} \right) - \frac{m}{\beta} \left(\frac{2}{\beta} - \frac{1}{\beta^2} - \frac{2}{\beta^3} - \frac{1}{\beta^4} \right) \frac{2a_1}{\beta} - \frac{3m + 1}{\beta^2}.$$

The right side of the inequality is strictly greater than zero for $a_1 \geq 2$. The minimum is therefore reached for either $a_1 = 0$, $a_1 = 1$ or $a_1 = 2$.

3) Difference Δ_{a_2} . The inspected difference is

$$\Delta_{a_2} = \frac{2a_2 + 1}{\beta^2} + c_2 + c_1 \frac{a_1}{\beta} + c_1 \frac{a_3}{\beta^2} + c_2 \frac{a_4}{\beta^2}.$$

For the difference $\Delta_{a_2} := Q_2(a_4, a_3, a_2 + 1, a_1, a_0) - Q_2(a_4, a_3, a_2, a_1, a_0)$ one needs to consider only $a_2 \leq m - 1$. Since $c_1, c_2 < 0$, we obtain

$$\Delta_{a_2} \leq \frac{2m - 1}{\beta^2} + c_2 = \frac{2m - 1}{\beta^2} - \frac{2m - 1}{\beta^2} - \frac{1}{\beta^4} = -\frac{1}{\beta^4} < 0.$$

The quadratic form is decreasing in the variable a_2 and the minimum is reached for the highest possible (w.r.t. Parry's condition) value of a_2 .

4) Difference Δ_{a_3} . The inspected difference is

$$\Delta_{a_3} = \frac{2a_3 + 1}{\beta^3} + c_3 + c_2 \frac{a_1}{\beta} + c_1 \frac{a_2}{\beta^2} + c_1 \frac{a_4}{\beta^3}.$$

We have shown that the minimum of the quadratic form is reached for $a_1 \in \{0, 1, 2\}$. It is suitable to discuss the cases $a_1 \in \{0, 1\}$ and $a_1 = 2$ separately.

Let us assume $a_1 \in \{0, 1\}$. Since $a_2, a_4 \leq m < \beta$ and $c_1, c_2 < 0$ we have

$$\Delta_{a_3} \geq \frac{1}{\beta^3} + c_3 + \frac{c_2}{\beta} + \frac{c_1}{\beta} + \frac{c_1}{\beta^2} = \frac{2(m-1)}{\beta^4} + \frac{1}{\beta^6} > 0.$$

This means that for $a_1 \in \{0, 1\}$ the minimum of the form is reached for $a_3 = 0$. Moreover, we already know that the minimum is reached for the highest possible value of a_2 (w.r.t. Parry's condition), therefore, we will inspect following candidates for the minimum $(a_4, 0, m-1, 1, 1)$ and $(a_4, 0, m, 0, 1)$.

The discussion for $a_1 = 2$ will be postponed for this moment.

5) Difference Δ_{a_4} . The inspected difference is

$$\Delta_{a_4} = \frac{2a_4 + 1}{\beta^4} + c_4 + c_3 \frac{a_1}{\beta} + c_2 \frac{a_2}{\beta^2} + c_1 \frac{a_3}{\beta^3}.$$

For the quintuple $(a_4, 0, m-1, 1, 1)$ we obtain

$$\Delta_{a_4} \geq \frac{1}{\beta^4} + c_4 + c_3 \frac{1}{\beta} + c_2 \frac{m-1}{\beta^2} > 0.$$

Hence we have the first candidate for the point where the form $Q_2(a_4, a_3, a_2, a_1, a_0)$ reaches its minimum

$$q_1 := (0, 0, m-1, 1, 1).$$

For the quintuple $(a_4, 0, m, 0, 1)$ we obtain

$$\Delta_{a_4} \leq \frac{2m+1}{\beta^4} + c_4 + c_2 \frac{m}{\beta^2} < 0.$$

Second candidate for the point where the form $Q_2(a_4, a_3, a_2, a_1, a_0)$ reaches its minimum is

$$q_2 := (m, 0, m, 0, 1).$$

Here, we get back to the case $a_1 = 2$. We inspect the difference Δ_{a_4} for the quintuple $(a_4, a_3, a_2, 2, 1)$. Note that due to the Parry's condition $a_2 \leq m-1$. We can estimate

$$\Delta_{a_4} \geq \frac{1}{\beta^4} + c_4 + c_3 \frac{2}{\beta} + c_2 \frac{m-1}{\beta^2} + c_1 \frac{m}{\beta^3} = \frac{2}{\beta^3} + \frac{2m-1}{\beta^4} - \frac{3}{\beta^5} > 0.$$

Hence we have to inspect the candidate $(0, a_3, a_2, 2, 1)$.

We treat separately the the case $a_3 = m$, which implies $a_2 = 0$. So we have the third candidate

$$q_3 := (0, m, 0, 2, 1)$$

and the cases $a_3 \leq m - 1$, where according to the negativity of Δ_{a_2} the value of corresponding variable is $a_2 = m - 1$. The difference Δ_{a_3} for the quintuple $(0, a_3, m - 1, 2, 1)$ is

$$\Delta_{a_3} = \frac{2a_3 + 1}{\beta^3} + c_3 + c_2 \frac{2}{\beta} + c_1 \frac{m - 1}{\beta^2} = \frac{2a_3 - 2m + 6}{\beta^3} + \frac{1 - m}{\beta^4} - \frac{2}{\beta^5} - \frac{1}{\beta^6}.$$

It is easy to see that for $a_3 \leq m - 3$ the $\Delta_{a_3} < 0$ and that for $a_3 \geq m - 2$ the $\Delta_{a_3} > 0$. Therefore the fourth candidate for the minimum is

$$q_4 := (0, m - 2, m - 1, 2, 1).$$

By computing the values of the quadratic form Q_2 in the points q_1, q_2, q_3 and q_4 we obtain

$$\frac{(\beta + 1)^2}{\beta^4} = Q_2(q_1) < \min\{Q_2(q_2), Q_2(q_3), Q_2(q_4)\}.$$

Therefore, $\text{Min}_5^2 = \beta^{-4}(\beta + 1)^2$ for $m \geq 3$.

Determining Max₃. Similarly as in the previous part of the proof we will now inspect the first differences of the quadratic form Q_1 in the individual variables.

From the definition of Max₃ we need to determine the maximal value of Q_1 over the set \mathcal{S}_1 , which allows the coefficients a_0, a_1, a_2 take any values in $\{0, 1, \dots, m\}$ with the admissibility condition. However, it is sufficient to consider $a_0 > 0$, since Q_1 reaches its maximum on such a point. Otherwise, we would have

$$\text{Max}_3 = |0 + a_1\beta' + a_2(\beta')^2| = |\beta'| |a_1 + a_2\beta'| \leq |\beta'| \text{Max}_2 < \text{Max}_2,$$

which is in contradiction with $\text{Max}_2 \leq \text{Max}_3$.

The inspection of the difference Δ_{a_0} of the quadratic form Q_1 is very similar to the one of Q_2 in the first part of the proof. We have

$$\Delta_{a_0} = 2a_0 + 1 + a_1c_1 + a_2c_2,$$

$a_0 \geq 1, c_1 < 0$ and $c_2 < 0$. Hence

$$\Delta_{a_0} \geq 3 + mc_1 + mc_2 \geq 3 - \frac{3m}{\beta} > 0,$$

the form is increasing in the variable a_0 and so the maximum is reached for the highest possible value of a_0 (w.r.t. Parry's condition).

Since

$$\Delta_{a_1} = \frac{2a_1 + 1}{\beta} + c_1a_0 + c_1 \frac{a_2}{\beta}$$

is a linear function in the variable a_1 and the coefficient $\frac{2}{\beta}$ at a_1 is positive, the difference Δ_{a_1} is either positive for all values of a_1 or negative for all values of a_1 or negative for some initial values of a_1 and then positive, regardless of the values of a_0 and a_2 .

Anyway, the maximum in the variable a_1 is reached for some extremal value of a_1 . The same reasoning works also for the maximum in the variable a_2 .

One can easily see that there are only six candidates fulfilling preceding conditions on the values of the variables a_2 , a_1 and a_0 :

	a_2	a_1	a_0
q_1	0	0	m
q_2	0	$m-1$	m
q_3	0	m	1
q_4	$m-1$	$m-1$	m
q_5	$m-1$	m	1
q_6	m	0	m

Finally, by computing the values of the quadratic form $Q_1(a_2, a_1, a_0)$ in the points q_1, \dots, q_6 we have

1. $Q_1(0, 0, m) = m^2$.
2. $Q_1(0, m-1, m) = m^2 + \frac{(m-1)^2}{\beta} - m(m-1) \left(\frac{1}{\beta} + \frac{1}{\beta^2} \right) < m^2$.
3. $Q_1(0, m, 1) = 1 + \frac{m^2}{\beta} - m \left(\frac{1}{\beta} + \frac{1}{\beta^2} \right) < m^2$.
4. $Q_1(m-1, m-1, m) = m^2 + \frac{(m-1)^2}{\beta} + \frac{(m-1)^2}{\beta^2} - m(m-1) \left(\frac{1}{\beta} + \frac{1}{\beta^2} \right) - (m-1)^2 \left(\frac{1}{\beta^2} + \frac{1}{\beta^3} \right) - m(m-1) \left(\frac{2}{\beta} - \frac{1}{\beta^2} - \frac{2}{\beta^3} - \frac{1}{\beta^4} \right) < m^2$.
5. $Q_1(m-1, m, 1) = 1 + \frac{m^2}{\beta} + \frac{(m-1)^2}{\beta^2} - m \left(\frac{1}{\beta} + \frac{1}{\beta^2} \right) - m(m-1) \left(\frac{1}{\beta^2} + \frac{1}{\beta^3} \right) - (m-1) \left(\frac{2}{\beta} - \frac{1}{\beta^2} - \frac{2}{\beta^3} - \frac{1}{\beta^4} \right) < m^2$.
6. $Q_1(m, 0, m) = m^2 + \frac{m^2}{\beta^2} - m^2 \left(\frac{2}{\beta} - \frac{1}{\beta^2} - \frac{2}{\beta^3} - \frac{1}{\beta^4} \right) < m^2$.

Therefore, $\text{Max}_3^2 = m^2$ and the proposition is proved.

5. Combinatorial properties of β -integers

It is known [19] that the set \mathbb{Z}_β has no accumulation points and that the distances between adjacent points in \mathbb{Z}_β take values in the set $\left\{ \sum_{i=1}^{\infty} \frac{t_i+k}{\beta^i} \mid k \in \mathbb{N}_0 \right\}$ provided that β is a Parry number. For β with the Rényi expansion $d_\beta(1) = m11$ the distances between adjacent points of \mathbb{Z}_β take three values, namely $1, \frac{1}{\beta} + \frac{1}{\beta^2}, \frac{1}{\beta}$. If we code these distances by letters a, b, c , respectively, the tiling of the positive real line by the non-negative elements of \mathbb{Z}_β can be represented as an infinite word $u_\beta := u_0 u_1 u_2 \dots$ in a ternary alphabet $\{a, b, c\}$. An example of the tiling of the real line by \mathbb{Z}_β for $m = 2$, and the corresponding infinite word u_β is shown on Figure 5.




Figure 1: The set $\mathbb{Z}_\beta \cap [0, +\infty)$ for $d_\beta(1) = 211$ and the corresponding word u_β .

The infinite word u_β for $d_\beta(1) = m11$ is invariant under the primitive substitution [10]

$$\begin{aligned} a &\mapsto a^m b, \\ b &\mapsto ac, \quad \text{where } a^m \text{ stands for } \underbrace{aa \cdots a}_{m \text{ times}}, \\ c &\mapsto a, \end{aligned} \tag{15}$$

For example, if $m = 2$, then we generate the fixed point of the substitution by

$$a \mapsto aab \mapsto aabaabac \mapsto aabaabacaabaabacaaba \mapsto \cdots$$

and we obtain the infinite word u_β from Figure 5.

For the description of combinatorial properties of an infinite word, crucial is the notion of a factor. We say that a finite word $w = w_0 w_1 \dots w_{n-1}$ is a factor of length n of the infinite word $u = u_0 u_1 u_2 \dots$, if for some i we have $w_0 w_1 \dots w_{n-1} = u_i u_{i+1} \dots u_{i+n-1}$. The function that to an integer n associates the number $\mathcal{C}(n)$ of different factors of length n in the infinite word u is called the complexity of u . It is known that the complexity of the infinite word u_β is $\mathcal{C}(n) = 2n + 1$ for all β such that $d_\beta(1) = m11$. A proof can be found in [11].


For $m = 1$, the infinite word u_β is called the Tribonacci word. It is known [7] that it is an Arnoux-Rauzy word, which means that for every length $n \in \mathbb{N}$ there exists a unique factor w of u_β of length n , such that aw , bw and cw are factors of u_β , and there exists a unique factor v of u_β of length n , such that va , vb and vc are factors of u_β .

It is known that if an infinite word is Arnoux-Rauzy, then the set of its factors is invariant under mirror image, i.e. with every factor $w_0 w_1 \dots w_{n-1}$ the infinite word contains the factor $w_{n-1} \dots w_1 w_0$. As a consequence, the infinite word u_β for $m \geq 2$ cannot be Arnoux-Rauzy, because its set of factors is not stable under mirror image. This can be seen easily from the substitution (15) under which u_β is invariant: $ca^m b$ is a factor of u_β , while $ba^m c$ is not.

Although the word u_β is for $m \geq 2$ not Arnoux-Rauzy, it is still true that for every length $n \in \mathbb{N}$ there exists a unique factor w of u_β of length n , such that aw , bw and cw are factors of u_β . However, one does not have a word v which could be extended with all the three letters to the right.

The invariance of the set of factors of the infinite word u_β decides about a property of the image of \mathbb{Z}_β under the field isomorphism $' : \mathbb{Q}(\beta) \mapsto \mathbb{Q}(\beta')$, the so-called Rauzy fractal. Figure 2 shows the set

$$\mathbb{Z}'_\beta = \left\{ \sum_{i=0}^k a_i \beta'^i \mid \sum_{i=0}^k a_i \beta^i \in \mathbb{Z}_\beta \right\}$$



2.1: Tribonacci numeration system.

2.2: Numeration system with minimal polynomial $x^3 - 2x^2 - x - 1$.

Figure 2: Central tiles in conjugated plane.

drawn in the complex plane, for the cases $m = 1$, $m = 2$. Bernat in [8] has shown that if $m = 1$, the closure of \mathbb{Z}'_β is a set centrally-symmetric with respect to the point $c = \frac{1}{2}(1 - \beta')^{-1}$. As it can be seen from Figure 2, \mathbb{Z}'_β for $m = 2$ does not have this property.

Acknowledgements

The authors acknowledge financial support of the Grant Agency of the Czech Republic GA ĀR 201/05/0169.

References

- [1] S. AKIYAMA. Cubic Pisot units with finite beta expansions. In: *Algebraic number theory and Diophantine analysis (Graz, 1998)*. de Gruyter, 2000, 11–26.
- [2] S. AKIYAMA. Self affine tiling and Pisot numeration system. In: K. GYÖRY, S. KANEMITSU (eds.), *Number theory and its applications (Kyoto, 1997)*. Vol. 2 of Dev. Math., Kluwer Acad. Publ., 1999, 7–17.
- [3] S. AKIYAMA, F. BASSINO, AND C. FROUGNY. Automata for arithmetic Meyer sets. In: *LATIN 2004: Theoretical informatics*. LNCS **2976**, Springer-Verlag, 2004, 252–261.
- [4] S. AKIYAMA, H. RAO, AND W. STEINER. A certain finiteness property of Pisot number systems. *J. Number Theory* **107** (2004), 135–160.
- [5] S. AKIYAMA AND T. SADAHIRO. A self-similar tiling generated by the minimal Pisot number. *Acta Math. Info. Univ. Ostraviensis* **6** (1998), 9–26.

- [6] P. AMBROŽ, C. FROUGNY, Z. MASÁKOVÁ, AND E. PELANTOVÁ. Arithmetics on number systems with irrational bases. *Bull. Belg. Math. Soc. Simon Stevin* **10** (2003), 641–659.
- [7] P. ARNOUX AND G. RAUZY. Représentation géométrique de suites de complexité $2n + 1$. *Bull. Soc. Math. France* **119** (1991), 199–215.
- [8] J. BERNAT. Computation of L_{\oplus} for Pisot numbers. Preprint IML Marseille, 2005.
- [9] Č. BURDÍK, C. FROUGNY, J.-P. GAZEAU, AND R. KREJCAR. Beta-integers as a group. In: *Dynamical systems (Luminy-Marseille, 1998)*. World Sci. Publishing, 2000, 125–136.
- [10] S. FABRE. Substitutions et β -systèmes de numération. *Theoret. Comput. Sci.* **137** (1995), 219–236.
- [11] C. FROUGNY, Z. MASÁKOVÁ, AND E. PELANTOVÁ. Complexity of infinite words associated with beta-expansions. *Theor. Inform. Appl.* **38** (2004), 163–185. Corrigendum. *Theor. Inform. Appl.* **38** (2004), 269–271.
- [12] C. FROUGNY AND B. SOLOMYAK. Finite beta expansions. *Ergod. Th. and Dynam. Sys.* **12** (1992), 713–723.
- [13] N. GJINI. A self-similar tiling generated by the Pisot number which is the root of the equation $x^3 - x^2 - 1 = 0$. *Osaka J. Math* **38** (2001), 303–319.
- [14] L.-S. GUIMOND, Z. MASÁKOVÁ, AND E. PELANTOVÁ. Arithmetics of beta-expansions. *Acta Arith.* **112** (2004), 23–40.
- [15] M. HOLLANDER. Linear numeration systems, finite beta-expansions, and discrete spectrum of substitution dynamical systems. PhD thesis, Washington University, 1996.
- [16] A. MESSAOUDI. Généralisation de la multiplication de Fibonacci. *Math. Slovaca* **50** (2000), 135–148.
- [17] A. MESSAOUDI. Tribonacci multiplication. *Appl. Math. Lett.* **15** (2002), 981–985.
- [18] W. PARRY. On the β -expansions of real numbers. *Acta Math. Acad. Sci. Hungar.* **11** (1960), 401–416.
- [19] W. P. THURSTON. Groups, tilings, and finite state automata. AMS Colloquium Lecture Notes, 1989.